

Complete Genome Sequence of *Bacillus amyloliquefaciens* Strain Co1-6, a Plant Growth-Promoting Rhizobacterium of *Calendula officinalis*

Martina Köberl,^{a,b} Richard A. White III,^b Sabine Erschen,^a Nora Spanberger,^{a,c} Tarek F. El-Arabi,^{d,e} Janet K. Jansson,^b Gabriele Berg^a

Graz University of Technology, Institute of Environmental Biotechnology, Graz, Austria^a; Pacific Northwest National Laboratory, Biological Sciences Division, Richland, Washington, USA^b; University of Graz, Institute of Plant Sciences, Graz, Austria^c; Ain Shams University, Faculty of Agriculture, Cairo, Egypt^d; Heliopolis University, Biotechnology Laboratory, Cairo, Egypt^e

The genome sequence of *Bacillus amyloliquefaciens* strain Co1-6, a plant growth-promoting rhizobacterium (PGPR) with broad-spectrum antagonistic activity against plant-pathogenic fungi, bacteria, and nematodes, consists of a single 3.9-Mb circular chromosome. The genome reveals genes putatively responsible for its promising biocontrol and PGP properties.

Received 24 June 2015 Accepted 9 July 2015 Published 13 August 2015

Citation Köberl M, White RA, III, Erschen S, Spanberger N, El-Arabi TF, Jansson JK, Berg G. 2015. Complete genome sequence of *Bacillus amyloliquefaciens* strain Co1-6, a plant growth-promoting rhizobacterium of *Calendula officinalis*. *Genome Announc* 3(4):e00862-15. doi:10.1128/genomeA.00862-15.

Copyright © 2015 Köberl et al. This is an open-access article distributed under the terms of the [Creative Commons Attribution 3.0 Unported license](https://creativecommons.org/licenses/by/3.0/).

Address correspondence to Martina Köberl, martina.koeberl@tugraz.at.

Bacillus amyloliquefaciens Co1-6 was isolated in October 2009 from the rhizosphere of the pot marigold *Calendula officinalis* L., cultivated on the organically managed Sekem farms in the northeastern desert region of Egypt (30°22'88"N 31°39'41"E) (1). The soil texture at the desert farm was classified as loamy sand, with a clay content of 4%, organic carbon content of 0.8%, and alkaline pH of 8.4 (2). Co1-6 was selected as a broad-spectrum antagonist exhibiting antifungal (*Verticillium dahliae*, *Rhizoctonia solani*, and *Fusarium culmorum*), antibacterial (*Ralstonia solanacearum*), and nematocidal (*Meloidogyne incognita*) activity against soilborne phytopathogens (3). For strains of the same population, induced systemic resistance of the host plant was identified as the major reason for their nematocidal activity (4). Treatment of chamomile plants (*Matricaria chamomilla*) with Co1-6 under field conditions resulted in elevated flavonoid contents of the blossoms (5).

Genomic DNA was extracted using the MasterPure DNA purification kit (Epicentre, Madison, WI, USA), modified with additional cell disruption steps comprising mechanical shredding with glass beads in a FastPrep instrument (MP Biomedicals, Santa Ana, CA, USA) and lysozyme-based cell wall digestion. PacBio RS libraries with inserts of 8 to 20 kb were constructed and sequenced at GATC Biotech (Konstanz, Germany).

Whole-genome shotgun sequencing yielded 245,374 raw reads with 1,486,144,876 bp of raw sequence. Assembly was completed with the Hierarchical Genome Assembly Process (HGAP) algorithm implemented in the PacBio SMRT Analysis software (Pacific Biosciences, Menlo Park, CA, USA) and resulted in a single circular chromosome of 3,922,431 bp, with 378.9-fold overall coverage and a G+C content of 46.85%.

The closest relative of Co1-6 based on the full-length 16S rRNA gene sequence is *B. amyloliquefaciens* subsp. *plantarum* FZB42 (NCBI reference sequence no. NR_075005, 99% sequence similarity). FZB42 is a well-known PGPR serving as the basis of a commercially available product (RhizoVital 42; ABiTEP GmbH,

Berlin, Germany) with the ability to stimulate plant growth and suppress plant pathogens (6). Digital DNA-DNA hybridization (DDH) using GGDC 2.0 (7–9) against the genome sequence of FZB42 (accession no. NC_009725) estimated a DDH of 80.30% ± 2.77%, indicating that they have 90.8% probability of being the same species but only 48.3% probability of being the same subspecies.

Annotation was conducted on the RAST Web server using RAST gene calling based on FIGfam version Release70 (10, 11), and additional annotation was completed on the BASys Web server using Glimmer gene prediction (12, 13). The genome annotation contained 3,913 predicted protein-coding genes, 86 tRNA and 19 rRNA loci, and 457 predicted SEED subsystem features.

The genome encodes synthases for mycosubtilin, plipastatin, and surfactin antibiotics, which most probably contribute to the promising abilities of Co1-6 for pathogen suppression. Co1-6 revealed six additional polyketide synthases, some at up to seven copies, and a dimodular nonribosomal peptide synthase. We further identified genes most probably involved in the direct promotion of plant growth, such as biosynthesis gene clusters for rhizobactin siderophores, spermidine, and auxin.

Nucleotide sequence accession numbers. This whole-genome shotgun project has been deposited in the European Nucleotide Archive under the accession no. [CVPA00000000](https://www.ebi.ac.uk/ena/submit/). The version described in this paper is the first version, CVPA01000000.

ACKNOWLEDGMENTS

This study was supported by the European Funds for Regional Development (EFRE), cofunded by the regional government of Styria (A3-11.P-33/2011-6), the Austrian Science Fund FWF (J 3638), cofunded by the European Commission, and the EU-Egypt Innovation Fund (RDI ENPI/2014/342-707).

A portion of this research is part of the Microbiomes in Transition Initiative at Pacific Northwest National Laboratory (PNNL). It was con-

ducted under the Laboratory-Directed Research and Development Program at PNNL, a multiprogram national laboratory operated by Battelle for the U.S. Department of Energy under contract DE-AC05-76RL01830.

REFERENCES

1. Köberl M, Müller H, Ramadan EM, Berg G. 2011. Desert farming benefits from microbial potential in arid soils and promotes diversity and plant health. *PLoS One* 6:e24452. <http://dx.doi.org/10.1371/journal.pone.0024452>.
2. Luske B, van der Kamp J. 2009. Carbon sequestration potential of reclaimed desert soils in Egypt. Louis Bolk Instituut & Soil and More International, Brussels, Belgium. <http://orgprints.org/16438/1/2192.pdf>.
3. Köberl M, Ramadan EM, Adam M, Cardinale M, Hallmann J, Heuer H, Smalla K, Berg G. 2013. *Bacillus* and *Streptomyces* were selected as broad-spectrum antagonists against soilborne pathogens from arid areas in Egypt. *FEMS Microbiol Lett* 342:168–178. <http://dx.doi.org/10.1111/1574-6968.12089>.
4. Adam M, Heuer H, Hallmann J. 2014. Bacterial antagonists of fungal pathogens also control root-knot nematodes by induced systemic resistance of tomato plants. *PLoS One* 9:e90402. <http://dx.doi.org/10.1371/journal.pone.0090402>.
5. Schmidt R, Köberl M, Mostafa A, Ramadan EM, Monschein M, Jensen KB, Bauer R, Berg G. 2014. Effects of bacterial inoculants on the indigenous microbiome and secondary metabolites of chamomile plants. *Front Microbiol* 5:64. <http://dx.doi.org/10.3389/fmicb.2014.00064>.
6. Chen XH, Koumoutsis A, Scholz R, Eisenreich A, Schneider K, Heine-meyer I, Morgenstern B, Voss B, Hess WR, Reva O, Junge H, Voigt B, Jungblut PR, Vater J, Süßmuth R, Liesegang H, Strittmatter A, Gottschalk G, Borriss R. 2007. Comparative analysis of the complete genome sequence of the plant growth-promoting bacterium *Bacillus amy-loliquefaciens* FZB42. *Nat Biotechnol* 25:1007–1014. <http://dx.doi.org/10.1038/nbt1325>.
7. Auch AF, von Jan M, Klenk HP, Göker M. 2010. Digital DNA-DNA hybridization for microbial species delineation by means of genome-to-genome sequence comparison. *Stand Genomic Sci* 2:117–134. <http://dx.doi.org/10.4056/sigs.531120>.
8. Auch AF, Klenk HP, Göker M. 2010. Standard operating procedure for calculating genome-to-genome distances based on high-scoring segment pairs. *Stand Genomic Sci* 2:142–148. <http://dx.doi.org/10.4056/sigs.541628>.
9. Meier-Kolthoff JP, Auch AF, Klenk HP, Göker M. 2013. Genome sequence-based species delimitation with confidence intervals and improved distance functions. *BMC Bioinformatics* 14:60. <http://dx.doi.org/10.1186/1471-2105-14-60>.
10. Aziz RK, Bartels D, Best AA, DeJongh M, Disz T, Edwards RA, Formsma K, Gerdes S, Glass EM, Kubal M, Meyer F, Olsen GJ, Olson R, Osterman AL, Overbeek RA, McNeil LK, Paarmann D, Paczian T, Parrello B, Pusch GD, Reich C, Stevens R, Vassieva O, Vonstein V, Wilke A, Zagnitko O. 2008. The RAST server: Rapid Annotations using Subsystems Technology. *BMC Genomics* 9:75. <http://dx.doi.org/10.1186/1471-2164-9-75>.
11. Overbeek R, Olson R, Pusch GD, Olsen GJ, Davis JJ, Disz T, Edwards RA, Gerdes S, Parrello B, Shukla M, Vonstein V, Wattam AR, Xia F, Stevens R. 2014. The SEED and the Rapid Annotation of microbial genomes using Subsystems Technology (RAST). *Nucleic Acids Res* 42:D206–D214. <http://dx.doi.org/10.1093/nar/gkt1226>.
12. Van Domselaar GH, Stothard P, Shrivastava S, Cruz JA, Guo A, Dong X, Lu P, Szafron D, Greiner R, Wishart DS. 2005. BASys: a Web server for automated bacterial genome annotation. *Nucleic Acids Res* 33:W455–W459. <http://dx.doi.org/10.1093/nar/gki593>.
13. Delcher AL, Harmon D, Kasif S, White O, Salzberg SL. 1999. Improved microbial gene identification with Glimmer. *Nucleic Acids Res* 27:4636–4641. <http://dx.doi.org/10.1093/nar/27.23.4636>.